

HEDIS® Measurement Year 2022 At-A-GlanceBehavioral Health Measures

We value everything you do to deliver quality care to our members — your patients — to ensure they have a positive healthcare experience. There are several HEDIS® behavioral health measures that providers can directly impact related to follow-up care for mental illness or substance use disorders, medication adherence, and metabolic monitoring. That's why we've created this easy-to-use At-A-Glance Toolkit. It will give you the tools you need to meet, document, and code HEDIS® measures. Together, we can provide the care and services our members need to stay healthy while also improving our quality scores and Star Ratings. Please contact your Provider Relations Representative if you need more information or have any questions.



HEDIS® Measure **Coding and Services** Intent and Recommendations Follow-Up Care for Children ** CPT® Codes for Initiation, Measure Intent: **Prescribed ADHD Medication Continuation and Maintenance** Provides an opportunity to track medication use in patients and Phases: 96150-96154, 98960-(ADD) provide the appropriate follow-98962, 99078, 99201-99205, Measure Specifications: 99211-99215, 99217-99219, up care to monitor clinical The percentage of children symptoms and potential 99241-99245, 99341-99345, newly prescribed ADHD adverse events. 99347, 99348-99350, 99381medication (no claims for 120 33387, 99391-99397, 99401days prior) who had at least Complete a comprehensive 99404, 99411-99412, 99483, three follow-up care visits medical and psychiatric 99510 within a 10-month period, one exam, including checklists of which was within 30 days of **Important Reminder for** for rating ADHD symptoms **Continuation and Maintenance** when the first ADHD medication before diagnosing and was dispensed. Two rates are **Phases of Treatment:** prescribing. reported: When prescribing a new 1. Initiation Phase. The Only one of the two visits ADHD medication, limit to a (during days 31–300) may be an percentage of members 14- to ages 6-12 with a e-visit or virtual check-in. 21-day supply and schedule prescription dispensed for a follow-up visit before the (**CPT® codes 98969, 98971, ADHD medication who had 99421 - 99444, 99457, G0017, patient or legal guardian one follow-up visit with a G2010, G2012, G2061 - G2063) leave the office to assess practitioner with how the medication is prescribing authority during **ADHD Medications:** working and to address any the 30-day Initiation Phase. side effects. Dexmethylphenidate, **2.** Continuation and Lisdexamfetamine, Educate and encourage the Maintenance (C&M) Phase. Methamphetamine, patient and parent or The percentage of members Dextroamphetamine, guardian to attend follow-Mehylphenidate, Clonidine, ages 6-12 with an up appointments monthly ambulatory prescription Guanfacine, Abtomoxetime until the child's symptoms dispensed for ADHD have stabilized, then every medication who were on 3-6 months for continued the medication for at least assessment of learning and 210 days and who, in behavior. If parents or addition to the visit in the guardian cancel the Initiation Phase, had at least patient's appointment, be two follow-up visits with a sure to reschedule right practitioner within 270 days away to keep within (9 months) after the measurement time periods. Initiation Phase ended. Submit the correct billing codes and utilize telehealth **Applicable Lines of Business:** options if needed.

This document is an informational resource designed to assist licensed healthcare practitioners in caring for their patients. Healthcare practitioners should use their professional judgment in using the information provided. HEDIS® measures are not a substitute for the care provided by licensed healthcare practitioners and patients are urged to consult with their healthcare practitioner for appropriate treatment. HEDIS® is a registered trademark of the National Committee for Quality Assurance (NCQA).

Medicaid

Age Group: 6-12 years



		T
Exclusion(s): Members in	Before prescribing ADHD	
hospice or with a diagnosis of	medication to younger	
narcolepsy	Medicaid patients, please refer	
	to the Preferred Drug List (PDL)	
Measurement Period:	on the health plan's state-	
The 12-month window starting	specific website.	
March 1 of the year prior to the		
measurement year and ending		
the last calendar day of		
February of the measurement		
year.		



Antidepressant Medication Management (AMM)

*Applicable Foster Care Measure

Measure Specifications:

The percentage of members ages 18 and older who were newly treated with an antidepressant medication (no claims for a period of 105 days prior) with a diagnosis of major depression and who remained on an antidepressant medication treatment. Two rates are reported:

- Effective Acute Phase Treatment. Members who remained on an antidepressant medication for at least 84 days (12 weeks).
- Effective Continuation Phase Treatment. Members who remained on an antidepressant medication for at least 180 days (6 months).

Applicable Lines of Business:

Medicaid Medicare Marketplace

Age Group:

18 years and older

Exclusion(s):

Members who are in hospice or members who do not have a diagnosis of major depression

Measurement Period:

The 12-month window starting on May 1 of the year prior to the measurement year and ending on April 30 of the measurement year.

Intent and Recommendations

Measure Intent:

Provides an opportunity to track antidepressant use in patients and provide appropriate follow-up care to monitor clinical worsening and/or suicide risk.

- Before diagnosing a patient with major depression, complete a comprehensive medical exam, including lab testing, which may identify a metabolic cause of depression. Accurate diagnosis drives appropriate treatments and interventions. Rule out medical or mental disorders that can produce symptoms similar to depression.
- Manage patient's depression with a systematic approach for accurate assessment and diagnosis. Begin with a nationally recognized tool such as the Patient Health Questionnaire (PHQ-9) using the billing code 96127 in conjunction with the ICD-10 diagnosis code Z13.

Engaging with and educating patients is the key to medication compliance. Consider taking these steps:

- Discuss how to take antidepressants, how they work, their benefits, and how long to take them.
- Tell your patients how long they can expect to be on an antidepressant before they start feeling better.
- Stress the importance of taking the medication even if they begin feeling better.

Coding and Services

Antidepressant Medications:

Bupropion, Vilazodone, Vortioxetine, Isocarboxazid, Phenelzine, Selegiline, Tranylcypromine, Nefazodone, Trazodone, Amitriptylinechlordiazepoxide, Amitriptylineperphenazine, Fluoxetineolanzapine, Desvenlafaxine, Duloxetine, Levomilnacipran, Venlafaxine, Citalopram, Escitalopram, Fluoxetine, Fluvoxamine, Paroxetine, Sertraline, Maprotiline, Mirtazapine, Amitriptyline, Amoxapine, Clomipramine, Desipramine, Doxepin (>6 mg), Imipramine, Nortriptyline, Protriptyline, Trimipramine



- 4. Talk about common side effects, how long they may last, and how to manage them.
- 5. Let your patient know what to do if they have questions or concerns.
- 6. Monitor with scheduled follow-up appointments.
- 7. Ask the patient to consider a psychotherapy referral. This may increase the chances of staying on medication and decrease the likelihood of a recurrence.

Before prescribing antidepressant medication to your Medicaid patients, please refer to the Preferred Drug List (PDL) on the health plan's state-specific website.

Before prescribing antidepressant medication to your Medicare patients, please refer to the health plan's formulary.



Metabolic Monitoring for Children and Adolescents on Antipsychotics (APM)

*Applicable Foster Care Measure

Measure Specifications:

The percentage of children and adolescents ages 1-17 who had two or more antipsychotic prescriptions and had metabolic testing. Three rates are reported:

- The percentage of children and adolescents on antipsychotics who received blood glucose testing.
- The percentage of children and adolescents on antipsychotics who received cholesterol testing.
- The percentage of children and adolescents on antipsychotics who received blood glucose <u>and</u> cholesterol testing.

<u>Applicable Line of Business:</u> Medicaid

Age Group: 1-17 years

Exclusion: Members in hospice

<u>Measurement Period:</u> Jan. 1 through Dec. 31 of a given calendar year.

Intent and Recommendations

Measure Intent:

Due to the potential negative health consequences associated with children developing cardiometabolic side effects from an antipsychotic medication, it is important to both establish a baseline and continuously monitor metabolic indices to ensure appropriate management of side effects.

- Encourage the parents or legal guardians of your patients prescribed antipsychotic medication to schedule an annual HbA1c test or glucose test and a LDL test when two or more antipsychotic medications are prescribed.
- Follow up with your patient's parent or legal guardian to discuss and educate them on lab results and what they mean.
- To increase compliance, consider ordering the annual HbA1c or glucose, and LDL tests as standing orders.
- Encourage parents or legal guardians to sign release of information forms and coordinate care with your patients' other treating medical and behavioral health specialists.

Coding and Services

** CPT® Codes for Blood Glucose Tests: 80047, 80048, 80050, 80053, 80069, 82947, 82950, 82951

- ** CPT® Codes for HbA1C Tests: 83036, 83037, 3044F, 3045F, 3046F
- ** CPT® Codes for LDL-C Tests: 80061, 83700, 83701, 83704, 83721, 3048F, 3049F, 3050F
- ** CPT® Codes for Cholesterol Tests other than LDL: 82465, 83718, 84478

Antipsychotic Medications:

Aripiprazole, Asenapine, Brexpiprazole, Cariprazine, Clozapine, Haloperidol, lloperidone, Loxapine, Lurasidone, Molindone, Olanzapine, Paliperidone, Pimozide, Quetiapine, Risperidone, Ziprasidone, Chlorpromazine, Fluphenazine, Perphenazine, Thioridazine, Trifluoperazine, Thiothixene, Aripiprazole, Aripiprazole lauroxil, Fluphenazine decanoate, Haloperidol decanoate, Olanzapine, Paliperidone palmitate, Fluoxetine-olanzapine, Perphenazine-amitriptyline, Prochlorperazine



Use of First-Line Psychosocial Care for Children and Adolescents on Antipsychotics (APP)

*Applicable Foster Care Measure

Measure Specifications:

The percentage of children and adolescents ages 1-17 who had a new prescription for an antipsychotic medication (no claims for a period of 120 days prior) and had documentation of psychosocial care as first-line treatment (90 days prior through 30 days after the dispensing date).

Applicable Line of Business:

Medicaid

Age Group: 1-17 years

Exclusion(s):

Members in hospice and members for whom first-line antipsychotic medications may be clinically appropriate.

Measurement Period: Jan. 1 through Dec. 1 of a given calendar year.

Intent and Recommendations

Measure Intent:

Children without primary indication for an antipsychotic and who are not given the benefit of a trial of psychosocial treatment first may unnecessarily incur the risks associated with antipsychotic medications. Since psychosocial interventions are associated with better outcomes, underuse of these therapies may lead to poorer mental and physical health outcomes.

- Before ordering a new antipsychotic prescription for your patient, consider a referral for a psychosocial assessment or psychosocial treatment.
- Ensure that the patient's psychosocial assessment or psychosocial treatment is part of their treatment record before writing a new prescription for an antipsychotic medication.
- Utilize telehealth options if needed.

Before prescribing antipsychotic medication to your younger Medicaid patients, please refer to the Preferred Drug List (PDL) on the health plan's state-specific website.

Coding and Services

** CPT® Codes for Psychological Care: 90832 - 90834, 90836 -

90840, 90845 - 90847, 90849, 90853, 90875, 90876, 90880

HCPCS Codes for Psychological Care:

G0176, G0177, G0409 - G0411, H0004, H0035 - H0040, H2000, H2001, H2011 - H2014, H2017 -H2020, S0201, S9480, S9484, S9485 With or without a telehealth modifier: 95, GT

Antipsychotic Medications:

Aripiprazole, Asenapine, Brexpiprazole, Cariprazine, Clozapine, Haloperidol, Iloperidone, Loxapine, Lurasidone, Molindone, Olanzapine, Paliperidone, Pimozide, Quetiapine, Risperidone, Ziprasidone, Chlorpromazine, Fluphenazine, Perphenazine, Thioridazine, Trifluoperazine, Thiothixene, Aripiprazole, Aripiprazole lauroxil, Fluphenazine decanoate, Haloperidol decanoate, Olanzapine, Paliperidone palmitate, Fluoxetine-olanzapine, Perphenazine-amitriptyline, Prochlorperazine



Follow-Up After Emergency Department Visit for Substance

HEDIS® Measure

Measure Specifications:

Use (FUA)

The percentage of emergency department (ED) visits for members ages 13 and older with a principal diagnosis of substance use disorder or any diagnosis of drug overdose. Two rates are reported:

- The percentage of ED visits for which the member received a follow-up visit or a pharmacotherapy dispensing event within 30 days of the ED visit (31 total days).
- 2. The percentage of ED visits for which the member received a follow-up visit or a pharmacotherapy dispensing event within 7 days of the ED visit (8 total days).

Applicable Lines of Business:

Medicaid Medicare

Age Group: 13 years and older

Exclusion(s):

ED visits that resulted in any inpatient stay the day of or within 30 days.

Members in hospice.

Measurement Period: Jan. 1 through Dec. 1 of a given calendar year.

Intent and Recommendations

Measure Intent:

To ensure care coordination for members who are discharged from the ED following high-risk substance use events, since those individuals may be particularly vulnerable to losing contact with the healthcare system.

Follow-up visits that occur on the same day as the ED discharge count for compliance.

- Schedule the 7-day follow-up visit within 5 days to allow flexibility in rescheduling.
- If the member's appointment does not occur within the first 7 days following their ED visit, please schedule the appointment to occur within 30 days post-ED visit.
- Keep in mind that patients in an ED setting may have a medical and comorbid substance use disorder diagnosis, so communication with the patient's PCP or medical specialist may be helpful in getting the patient into substance use disorder treatment.
- Utilize telehealth options if needed.

Coding and Services

Visit Setting Unspecified **CPT®
Codes: 90791-90792, 90832-90834,
90836-90840, 90845, 90847,
90849, 90853, 90875-90876,
99221-99223, 99231-99233,
99238-99239, 99251-99255
Outpatient POS: 03, 05, 07, 09, 1120, 22, 33, 49, 50, 71-72

*ICD-10 AOD Abuse and
Dependence and Substance
Induced Disorder Codes:
F10.xx-F19.xx (excludes remission codes)

<u>Unintentional Drug Overdose</u> <u>Codes:</u> example T40.0X1A

BH Outpatient **CPT® Codes: 98960-98962, 99078, 99201-99205, 99211-99215, 99241-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411-99412, 99483, 99492-99494, 99510

BH Outpatient HCPCS: G0155, G0176-G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036-H0037, H0039-H0040, H2010-H2011, H2013-H2020, T1015 Partial Hospitalization POS: 52

Partial Hospitalization/Intensive Outpatient HCPCS Codes: G0410-G0411, H0035, H2001, H2012, S0201, S9480, S9484-S9485

<u>Substance Use Disorder Services</u> <u>HCPCS Codes:</u> G0396-G0397, G0443, H0001, H0005, H0007, H0015-H0016, H0022, H0047, H0050, H2035-H2036, T1006, T1012



HEDIS® Measure	Intent and Recommendations	Coding and Services
		Substance Use Disorder Services UB Rev Codes: 0906, 0944-0945
		Behavioral Health Assessment **CPT® Codes: 99408-99409
		Behavioral Health Assessment HCPCS Codes: G0396-G0397, G0442, G2011, H0001-H0002, H0031, H0049
		Substance Use Services HCPCS Codes: H0006, H0028
		AOD Medication Treatment HCPCS Codes: H0020, H0033, J0570-J0575, J2315, Q9991-Q9992, S0109
		OUD Weekly Drug Treatment Service HCPCS Codes: G2067-G2073



Follow-Up After Hospitalization for Mental Illness (FUH)

*Applicable Foster Care Measure

Measure Specifications:

The percentage of discharges for members 6 years of age and older who were hospitalized for treatment of selected mental illness or intentional self-harm diagnoses and who had a follow-up visit with a mental health provider. Two rates are reported:

- The percentage of discharges for which the member received follow-up within 30 days <u>after</u> discharge.
- The percentage of discharges for which the member received follow-up within 7 days <u>after</u> discharge.

Applicable Lines of Business:

Medicaid Medicare Marketplace

Age Group: 6 years and older

Exclusion(s):

Non-acute inpatient. Members in hospice.

Measurement Period: Jan. 1 through Dec. 1 of a given calendar year.

Intent and Recommendations

Measure Intent:

An outpatient visit with a mental health practitioner after discharge is recommended to make sure that the patient's transition to the home or work environment is supported and that gains made during hospitalization are not lost. It also helps health care providers detect early post-hospitalization reactions or medication problems and provide continuing care.

Follow-up visits that occur on the same day as the IP discharge do not count.

- Schedule the 7-day follow-up visit within 5 days of discharge to allow flexibility in rescheduling.
- If the member's appointment does not occur within the first 7 days post-discharge, please schedule the appointment to occur within 30 days.
- Engage with and educate the member and guardian on the importance of follow-up care
- Utilize telehealth options if needed.

Coding and Services

Visit Setting Unspecified **CPT®
Codes: 90791-90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875-90876, 99221-99223, 99231-99233, 99238-99239, 99251-99255

BH Outpatient **CPT® Codes: 98960-98962, 99078, 99201-99205, 99211-99215, 99241-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411-99412, 99483, 99492-99494, 99510

<u>Electroconvulsive Therapy **CPT®</u> Code: 90870

Observation CPT® Codes: 99217-99220

<u>Transitional Care Management</u> <u>Services **CPT® Codes:</u> 99495-99496

BH Outpatient HCPCS:

G0155, G0176-G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010-H2011, H2013-H2020, T1015

Community Mental Health Center POS: 53

ICD10-PCS Codes: GZB0ZZZ-GZB4ZZZ

Ambulatory Surgical Center POS: 24
Outpatient POS: 03, 05, 07, 09, 1120, 22, 33, 49, 50, 71-72
Partial Hospitalization POS: 52
Partial Hospitalization/Intensive
Outpatient HCPCS: G0410-G0411,
H0035, H2001, H2012, S0201,
S9480, S9484-S9485



		Complete neath.
HEDIS® Measure	Intent and Recommendations	Coding and Services
		Partial Hospitalization/Intensive Outpatient UB Rev: 0905, 0907, 0912, 0913
		Telehealth POS: 02
		Behavioral Healthcare Setting UB Rev Codes: 0513, 0900-0919
		Telephone Visits **CPT® Codes: 98966-98968, 99441-99443
		Psychiatric Collaborative Care Management **CPT® Codes: 99492-99494
		Psychiatric Collaborative Care Management HCPCS Code: G0512
		OUD Monthly Office-Based Treatment HCPCS Codes: G2086-G2087
		Observation CPT® Codes: 99217- 99220
		Residential Behavioral Health Treatment HCPCS Codes: H0017-H0019, T2048
		Telephone Visits CPT® Codes: 98966-98968, 98441-98443
		Online Assessments CPT® Codes: 98969, 98971-98972, 99421-99444, 99457
		AOD Medication Treatment HCPCS Codes: H0020, H0033, J0570-J0575, J2315, Q9991, Q9992, S0109
	I	1



Follow-Up After High-Intensity Care for Substance Use Disorder (FUI)

Measure Specifications:

The percentage of acute inpatient hospitalizations, residential treatments, or detoxification visits for a diagnosis of substance use disorder among members ages 13 and older that result in a follow-up visit or service for substance use disorder. Compliance includes:

- A follow-up visit or event with any practitioner for a principal diagnosis of substance use disorder within 30 days after the visit or discharge.
- A follow-up visit or event with any practitioner for a principal diagnosis of substance use disorder within 7 days after the visit or discharge.

Applicable Lines of Business:

Medicaid Medicare

Age Group: 13 years and older **Exclusion(s):** Members in hospice and non-acute inpatient

<u>Measurement Period:</u> Jan. 1 through Dec. 1 of a given calendar year.

Intent and Recommendations

Measure Intent:

Timely follow up and continuity of care following a high-intensity event for a diagnosis of substance use disorder is critical, as individuals receiving substance use disorder care in these settings are vulnerable to losing contact with the healthcare system.

Follow-up visits that occur on the same day as the discharge from IP, residential treatment, or detoxification visits do not count.

- Schedule the 7-day follow-up visit within 5 days to allow flexibility in rescheduling. Be sure to involve the patient's caregiver or patent/guardian in any follow-up instructions after the visit or discharge.
- If the patient's appointment does not occur within the first 7 days post-visit/discharge, please schedule the appointment to occur within 30 days post-visit/discharge.
- Follow-up does not include detoxification.
- Methadone is not included on the medication lists for this measure.
- Engage with and educate the patient and/or their parent or legal guardian on the importance of follow-up care
- Utilize telehealth options or phone visits if needed.

Coding and Services

*ICD-10 AOD Abuse and

Dependence and Substance
Induced Disorder Codes: F10.xx-F19.xx (excludes remission codes)

Inpatient Stay UB Rev Codes: 0100-0101, 0110-0114, 0116-0124, 0126-0134, 0136-0144, 0146-0154, 0156-0160, 0164, 0167, 0169-0174, 0179, 0190-0194, 0199-0204, 0206-0214, 1000-1002

Visit Setting Unspecified **CPT®
Codes: 90791-90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875-90876, 99221-99223, 99231-99233, 99238-99239, 99251-99255

Outpatient POS: 03, 05, 07, 09, 11-20, 22, 33, 49, 50, 71-72

BH Outpatient **CPT® Codes: 98960-98962, 99078, 99201-99205, 99211-99215, 99241-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411-99412, 99483, 99492-99494, 99510

BH Outpatient HCPCS:

G0155, G0176-G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010-H2011, H2013-H2020, T1015

Partial Hospitalization POS: 52

Partial Hospitalization/Intensive Outpatient HCPCS Codes: G0410-G0411, H0035, H2001, H2012, S0201, S9480, S9484-S9485

Non-residential Substance Abuse Treatment Facility POS: 57-58



HEDIS® Measure	Intent and Recommendations	Coding and Services
		OUD Weekly Drug Treatment Service HCPCS Codes: G2067-G2073
		Pharmacotherapy dispensing event Opioid Use Disorder Treatment Medications: Naltrexone (oral and injectable), Buprenorphine (sublingual tablet, injection, implant) 1, Buprenorphine/naloxone (sublingual tablet, buccal film, sublingual film) Alcohol Use Disorder Treatment Medications: Disulfiram (oral), Naltrexone (oral and injectable), Acamprosate (oral and delayed-release tablet)
		¹ Buprenorphine administered via transdermal patch or buccal film are not included because they are FDA-approved for the treatment of pain and not for opioid use disorder.



Follow-Up after Emergency Department Visit for Mental Illness (FUM)

*Applicable Foster Care Measure

Measure Specifications:

The percentage of emergency department (ED) visits for members ages 6 and older with a principal diagnosis of mental illness or intentional self-harm who had a follow-up visit for mental illness. Two rates are reported:

- The percentage of ED visits for which the member received follow-up care within 30 days of the ED visit (31 total days).
- The percentage of ED visits for which the member received follow-up care within 7 days of the ED visit (8 total days).

Applicable Lines of Business:

Medicaid Medicare

Age Group: 6 years and older Exclusion(s): ED visits that resulted in any inpatient stay the day of or within 30 days.
Members in hospice.

<u>Measurement Period:</u> Jan. 1 through Dec. 1 of a given calendar year.

Intent and Recommendations

Measure Intent:

Research suggests that for people with serious mental illness, both low-intensity interventions, such as appointment reminders, and high-intensity interventions, such as assertive community treatment, can be effective following an ED visit. These interventions encourage follow-up care in the outpatient setting.

Follow-up visits can occur on the same day as the ED discharge

- Schedule the 7-day follow-up visit within 5 days to allow flexibility in rescheduling.
- If the member's appointment does not occur within the first 7 days post-ED visit, please schedule the appointment to occur within 30 days post-ED visit.
- Keep in mind that patients in an ED setting may have a medical and comorbid mental health diagnosis, so communication with the patient's PCP or medical specialist may be helpful in getting the patient into treatment for their mental illness.
- Utilize telehealth options if needed.

The follow-up visit must have a principal diagnosis of a mental health disorder or intentional self-harm.

Coding and Services

*ICD-10 Mental Illness Diagnosis Codes: F03.9x, F20-25.xx, F28-34. xx, F39-45.xx, F48.xx, F50-53.xx, F59-60.xx, F63-66.xx, F68-69.xx, F80-82.xx, F84.xx, F88-93.xx, F95.xx, F98-99.xx

* ICD-10 Intentional Self-Harm Diagnosis Codes example: T39.92XA

<u>Visit Setting Unspecified **CPT®</u>
<u>Codes:</u> 90791-90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875-90876, 99221-99223, 99231-99233, 99238-99239, 99251-99255

BH Outpatient **CPT® Codes: 98960-98962, 99078, 99201-99205, 99211-99215, 99241-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411-99412, 99483, 99492-99494, 99510

BH Outpatient HCPCS:

G0155, G0176-G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010-H2011, H2013-H2020, T1015

Partial Hospitalization POS: 52

Partial Hospitalization/Intensive Outpatient HCPCS Codes: G0410-G0411, H0035, H2001, H2012, S0201, S9480, S9484-S9485

<u>Community Mental Health Center</u> POS: 53

Electroconvulsive Therapy **CPT Codes: 90870



HEDIS® Measure	Intent and Recommendations	Coding and Services
		Ambulatory Surgical Center POS: 24 Outpatient POS: 03, 05, 07, 09, 11- 20, 22, 33, 49, 50, 71-72
		Telehealth POS: 02
		Observation CPT® Codes: 99217- 99220
		Telephone Visits **CPT® Codes: 98966-98968, 99441-99443
		Behavioral Healthcare Setting UB Rev Codes: 0513, 0900-0919



Initiation and Engagement of Substance Use Disorder Treatment (IET)

HEDIS® Measure

Measure Specifications:

The percentage members ages 13 and older with a <u>new</u> episode of substance use disorder (<u>and</u> no substance use disorder diagnoses within the past 194 days) who received the following.:

- Initiation of Substance Use
 Disorder Treatment. The
 percentage of members who
 initiate treatment through an
 inpatient substance use
 disorder admission, outpatient
 visit, intensive outpatient
 encounter or partial
 hospitalization, telehealth, or
 medication treatment within
 14 days of the diagnosis.
- Engagement of Substance Use
 Disorder Treatment. The
 percentage of members who
 initiated treatment and who
 were engaged in ongoing
 substance use disorder
 treatment within 34 days of the
 initiation visit.

Applicable Lines of Business:

Medicaid Medicare Marketplace

Age Group: 13 years and older

Exclusion(s):

Members in hospice or substance use disorder episodes that occurred during the 194 days prior to the new substance use disorder episode date.

Intent and Recommendations

Measure Intent:

Individuals who engage in early substance use disorder treatment have been found to have decreased odds of negative outcomes, including mortality. The intent of this measure is to measure access to evidence-based substance use disorder treatment for patients beginning a new episode of treatment.

- A PCP or medical specialist, along with BH practitioners and providers, may provide the substance use disorder diagnosis in a variety of settings, such as, but not limited to, a medical ED visit, PCP office visit, acute IP medical treatment, or treatment for detox. Once the patient has a new substance use disorder diagnosis, the initiation phase begins.
- Schedule the initial 14-day follow-up visit within 10 days of a new substance use disorder diagnosis to allow flexibility in rescheduling.
- Utilize telehealth options if needed.

At the end of the initial follow-up appointment, schedule two more appointments to occur within 34 days of the initial visit.

Coding and Services

*ICD-10 AOD Abuse and

Dependence and Substance
Induced Disorder Codes: F10.xxF19.xx (excludes remission codes)

Visit Setting Unspecified **CPT®
Codes: 90791-90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875-90876, 99221-99223, 99231-99233, 99238-99239, 99251-99255

Outpatient POS: 03, 05, 07, 09, 11-20, 22, 33, 49, 50, 71-72

BH Outpatient **CPT® Codes: 98960-98962, 99078, 99201-99205, 99211-99215, 99241-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411-99412, 99483, 99492-99494, 99510

BH Outpatient HCPCS:

G0155, G0176-G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010-H2011, H2013-H2020, T1015

Partial Hospitalization POS: 52

Partial Hospitalization/Intensive Outpatient HCPCS Codes: G0410-G0411, H0035, H2001, H2012, S0201, S9480, S9484-S9485

Non-residential Substance Abuse Treatment Facility POS: 57-58

Community Mental Health Center POS: 53

Telehealth POS: 02



HEDIS® Measure	Intent and Recommendations	Coding and Services
Measurement Period: New episodes of substance use disorder, Nov. 15 of the year prior to the measurement year through Nov. 14 of the measurement year.	 When treating a member for issues related to substance use disorder, remember to code the diagnosis on every claim. Encourage patients and their caregivers or parents/guardians to sign the appropriate release of information forms and coordinate care with their medical and behavioral health providers. Follow-up care does not include detoxification. Exclude all detoxification events (HCPCS H0008-H0014, ICD-10 PCS HZ2ZZZZ, UB Rev 0116, 0126, 0136, 0146, 0156) when identifying follow-up visits for numerator compliance. 	Substance Use Disorder Services HCPCS Codes: G0396-G0397, G0443, H0001, H0005, H0007, H0015-H0016, H0022, H0047, H0050, H2035-H2036, T1006, T1012 OUD Weekly Non-Drug Treatment Service HCPCS Codes: G2071, G2074-G2077, G2080 OUD Monthly Office-Based Treatment HCPCS Codes: G2086-G2087 Observation CPT® Codes: 99217-99220 Residential Behavioral Health Treatment HCPCS Codes: H0017-H0019, T2048 Telephone Visits CPT® Codes: 98966-98968, 98441-98443 Online Assessments CPT® Codes: 98969, 98971-98972, 99421- 99444, 99457 AOD Medication Treatment HCPCS Codes: H0020, H0033, J0570-J0575, J2315, Q9991, Q9992, S0109 OUD Weekly Drug Treatment Service HCPCS Codes: G2067-G2073



HEDIS® Measure	Intent and Recommendations	Coding and Services
		Pharmacotherapy dispensing event Opioid Use Disorder Treatment Medications: Naltrexone (oral and injectable), Buprenorphine (sublingual tablet, injection, implant) 1, Buprenorphine/naloxone (sublingual tablet, buccal film, sublingual film) Alcohol Use Disorder Treatment Medications: Disulfiram (oral), Naltrexone (oral and injectable), Acamprosate (oral and delayed-release tablet)
		¹ Buprenorphine administered via transdermal patch or buccal film are not included because they are FDA-approved for the treatment of pain and not for opioid use disorder.



HEDIS® Measure **Intent and Recommendations Coding and Services Diabetes Screening for People Measure Intent:** *ICD-10 Codes for Schizophrenia: with Schizophrenia or Bipolar Diabetes screening for individuals F20.0-F20.5, F20.81, F20.89, F20.9, Disorder who are Using with schizophrenia or bipolar F25.0-F25.1, F25.8-F25.9 **Antipsychotic Medications (SSD)** disorder who are prescribed an *ICD-10 Codes for Bipolar antipsychotic medication may lead Disorder: F30.10-F30.13, F30.2-**Measure Specifications:** to earlier identification and F30.4, F30.8-F30.9, F31.10-F31.13, The percentage of members ages treatment of diabetes. A glucose 18-64 with schizophrenia, F31.2, F31.30-F31.32, F31.4-F31.5, test or an HbA1c test performed F31.60-F31.64, F31.70-F31.78 schizoaffective disorder, or during the measurement year will bipolar disorder who were close this gap. **CPT® Codes for Glucose Tests: dispensed an antipsychotic 80047, 80048, 80050, 80053, • Encourage members with medication and had a diabetes 80069, 82947, 82950, 82951 schizophrenia or bipolar screening test during the disorders who are also on **CPT® Codes for HbA1c Tests: measurement year. antipsychotic medication to 83036, 83037 CPT II: 3044F, **Applicable Lines of Business:** schedule an annual HbA1c test 3045F, 3046F, ≥7%-<8%: 3051F, or glucose test at the time of Medicaid ≥8%-≤9%: 3052F their visit. **Age Group:** 18-64 years • To increase compliance, consider ordering the HbA1c or Exclusion(s): glucose test as an annual Members diagnosed with standing order. diabetes. • BH providers (MD, NP or other professional with lab ordering Members diagnosed with ability) can order diabetic tests schizophrenia or bipolar but who for patients who do not have did not receive antipsychotic regular contact with their PCP. medication. The BH provider then Members in hospice. coordinates medical management with the PCP. Measurement Period: Jan. 1 through Dec. 31 of a given calendar year.