

Clinical Policy: Pancreas Transplantation

Reference Number: NC.CP.MP.102

Date of Last Revision: 04/22

[Coding Implications](#)
[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

This policy describes the medical necessity requirements for pancreas transplantation procedures. Multiple types of pancreas transplants are effective therapeutic options for arresting the progression of complications of diabetes mellitus and improving the quality of life for diabetic patients, including simultaneous pancreas kidney transplant (SPK), pancreas after kidney transplant (PAK), pancreas transplant alone (PTA), and islet cell transplant.¹

Policy/Criteria

- I. It is the policy of Carolina Complete Health that a pancreas transplant is **medically necessary** when meeting all of the following:
 - A. Severely disabling and potentially life-threatening complications due to hypoglycemic unawareness and labile insulin-dependent diabetes (Type I DM) that persists in spite of optimal medical management;
 - B. Does not have ANY of the following contraindications:
 1. Malignancy with high risk of recurrence or death related to cancer;
 2. Glomerular filtration rate $< 40 \text{ mL}/\text{min}/1.73\text{m}^2$ unless being considered for multi-organ transplant;
 3. Stroke, acute coronary syndrome, or myocardial infarction (excluding demand ischemia) within 30 days;
 4. Acute liver failure, or cirrhosis with portal hypertension or synthetic dysfunction unless being considered for multi-organ transplant;
 5. Acute renal failure with rising creatinine or on dialysis and low likelihood of recovery;
 6. Septic shock;
 7. Chronic infection with highly virulent and/or resistant microbes that are poorly controlled pre-transplant;
 8. Active tuberculosis infection;
 9. HIV infection with detectable viral load;
 10. Progressive cognitive impairment;
 11. Inability to adhere to the regimen necessary to preserve the transplant, even with caregiver support;
 12. Active substance use or dependence including current tobacco use, vaping, marijuana smoking, or IV drug use without convincing evidence of risk reduction behaviors, such as meaningful and/or long-term participation in therapy for substance abuse and/or dependence. Serial blood and urine testing may be used to verify abstinence from substances that are of concern;
 13. Other severe uncontrolled medical condition expected to limit survival after transplant;

CLINICAL POLICY Pancreas Transplantation

C. Request is for one of the following procedures and meets the corresponding criteria:

1. *PTA*, meets all:
 - a. Recurrent, severe, and potentially life-threatening metabolic complications that require medical attention, as documented by chart notes, emergency room visits, or hospitalizations, including any of the following: i. Severe hypoglycemia unawareness; ii. Marked hyperglycemia; iii. Recurring severe ketoacidosis;
 - b. Clinical and emotional problems with exogenous insulin therapy that are so severe as to be incapacitating or consistent failure of insulin-based management to prevent acute complications;
2. *SPK* with end-stage renal disease (ESRD), as evidenced by uremia;
3. *PAK transplant*, meets all:
 - a. Underwent successful kidney transplant without significant chronic rejection of kidney transplant;
 - b. Stable kidney transplant function, as defined by both:
 - i. Stable creatinine clearance ≥ 30 mL/min,
 - ii. Absence of significant proteinuria.

II. It is the policy of Carolina Complete Health that *autologous islet cell transplants* are considered **medically necessary** as an adjunct procedure to a total or near total pancreatectomy for severe, refractory pancreatitis.

III. It is the policy of Carolina Complete Health that *pancreas re-transplantation* are considered **medically necessary** after one failed primary pancreas transplant.

IV. It is the policy of Carolina Complete Health that current evidence does not support the use of pancreas transplant procedures for any of the following indications:

- a. Re-transplantations after two or more failed primary pancreas transplantations;
- b. Allogeneic islet cell transplantation or xenotransplantation;
- c. SPK transplantation for patients with amputation due to peripheral obstructive vascular disease;
- d. For the treatment of all other conditions than those specified above.

Background

The American Diabetes Association³ defines diabetes mellitus as a group of metabolic diseases characterized by hyperglycemia that results from defects in insulin secretion, insulin action, or both. According to the Centers for Disease Control and Prevention⁴ estimations in 2020, approximately 34.2 million people or 10.5% of the U.S. population have diabetes with approximately 7.3 million undiagnosed cases. Chronic hyperglycemia existing in diabetic patients facilitates long term organ damage, especially to the eyes, kidneys, nerves, and blood vessels.³

The prevalent type 2 diabetes is caused by a resistance to insulin action and an inadequate compensatory insulin secretory response.³ Type 1 diabetes is caused by immune mediated

CLINICAL POLICY Pancreas Transplantation

destruction of the insulin secreting pancreatic β cells.⁵ Islet cell autoantibodies, insulin autoantibodies, autoantibodies to glutamic acid decarboxylase, zinc transporter 8 (ZnT8A), and autoantibodies to the tyrosine phosphatase IA-2 and IA-2 β are serological markers of the pancreatic β cell destruction observed in type 1 diabetes.^{3,5-6}

Pancreas transplantation allows for the possibility to restore glucose regulated endogenous secretion, decrease the progression of diabetic complications, and improve quality of life in patients with diabetes.^{1,7} Pancreas transplantation is the only proven method to restore normoglycemia in type 1 diabetic patients.⁸ Simultaneous pancreas kidney transplant (SPK), pancreas after kidney transplant (PAK), and pancreas transplant alone (PTA) are primarily performed on patients with type 1 diabetes.⁸ SPK is an established procedure for diabetic patients with advanced chronic kidney disease or end stage kidney disease and accounts for approximately 90% of pancreas transplants performed in the United States.⁹

A 2011 study by Gruessner¹⁰ reviewed the outcomes of SPK, PAK, and PTA transplantations according to follow-up data collected by the International Pancreas Transplant Registry. Patient survival rates were reported to be over 95% after 1 year and over 83% at 5 years post-transplant. The highest graft survival rates were observed in SPK transplants at 86% for pancreas and 93% for kidney graft function 1-year post-transplant. PAK procedures displayed graft function at 80%, while PTA had graft function at 78% one year after transplantation.¹⁰ Graft survival rate is defined as total freedom from insulin therapy, normal fasting blood glucose concentrations, and normal or only slightly elevated hemoglobin A1C values.¹¹ The study demonstrated that pancreas transplantation offers excellent outcomes for patients with labile diabetes due to the improvement in patient survival and graft function shown in all 3 categories over the course of 24 years.¹⁰

Patients undergoing pancreas transplantation, especially SPK transplant, require extensive immunosuppression regimens.¹ It is theorized that pancreas transplant recipients require higher levels of immunosuppression therapy than other solid organ transplants due to the immunogenicity of the pancreas or the autoimmune status of the recipient.¹²

During pancreatic islet autotransplantation, Islet β cells are transferred into the liver through the portal vein of the recipient.¹ Pancreatic islet autotransplantation is performed following a pancreatectomy in patients with severe chronic pancreatitis. Chronic pancreatitis is a debilitating disease which causes diarrhea, weight loss, poor quality of life, and severe abdominal pain that is difficult to alleviate with pharmacological treatment or other therapeutic measures.^{1,13} Due to the excessive pain observed in patients with chronic pancreatitis, pain control is a primary goal of pancreatectomy and pancreatic islet autotransplantation.¹³

Coding Implications

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2019, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not

CLINICAL POLICY Pancreas Transplantation

guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

CPT®* Codes	Description
48160	Pancreatectomy, total or subtotal, with autologous transplantation of pancreas of pancreatic islet cells
48550	Donor pancreatectomy (including cold preservation), with or without duodenal segment for transplantation
48551	Backbench standard preparation of cadaver donor pancreas allograft prior to transplantation, including dissection of allograft from surrounding soft tissues, splenectomy, duodenotomy, ligation of bile duct, ligation of mesenteric vessels, and Y-graft arterial anastomoses from iliac artery to superior mesenteric artery and to splenic artery
48552	Backbench reconstruction of cadaver donor pancreas allograft prior to transplantation, venous anastomosis, each
48554	Transplantation of pancreatic allograft
48556	Removal of transplanted pancreatic allograft
50300	Donor nephrectomy (including cold preservation) from cadaver donor, unilateral or bilateral
50320	Donor nephrectomy (including cold preservation); open, from living donor
50323	Backbench standard preparation of cadaver donor renal allograft prior to transplantation, including dissection and removal of perinephric fat, diaphragmatic and retroperitoneal attachments, excision of adrenal gland, and preparation of ureter(s), renal vein(s), and renal artery(s), ligating branches, as necessary
50325	Backbench standard preparation of living donor renal allograft (open or laparoscopic) prior to transplantation, including dissection and removal of perinephric fat and preparation of ureter(s), renal vein(s), and renal artery(s), ligating branches, as necessary
50327	Backbench reconstruction of cadaver or living donor renal allograft prior to transplantation; venous anastomosis, each
50340	Recipient nephrectomy (separate procedure)
50360	Renal allotransplantation, implantation of graft; without recipient nephrectomy
50365	Renal allotransplantation, implantation of graft; with recipient nephrectomy

HCPCS Codes	Description
S2065	Simultaneous pancreas kidney transplantation

ICD-10 Diagnosis Codes that Support Coverage Criteria

+ Indicates a code requiring an additional character

ICD-10-CM Code	Description
E10.10-E10.9	Type 1 diabetes mellitus with kidney complications
K86.0	Alcohol-induced chronic pancreatitis

CLINICAL POLICY Pancreas Transplantation

K86.1	Other chronic pancreatitis
N18.4	Chronic kidney disease, stage 4 (severe)
N18.5	Chronic kidney disease, stage 5
N18.6	End stage renal disease
ICD-10-CM Code	Description
Z94.0	Kidney transplant status

Reviews, Revisions, and Approvals	Revision Date	Approval Date
Policy adapted for use by North Carolina Health Plan (Carolina Complete Health), per state feedback and requirements: replaced requirement for failure or non-existence of medical treatment for condition and indication of DM 1 and definitions of insulin dependence per specific C peptide values/C peptide values and BMI with the indication of severe complications due to hypoglycemic unawareness and labile Type 1 DM. Replaced contraindications of malignancy in past 2 years or metastasized malignancy with “active, potentially life-threatening malignancy.”	06/19	6/19
Annual review completed. No changes to policy content.	03/21	03/21
Section I.B, removed contraindication of “severely limited functional status with poor rehabilitation potential.” Replaced “Psychiatric or psychological condition associated with the inability to cooperate or comply with medical therapy” and the contraindication regarding non-compliance with medical therapy with “Inability to adhere to the regimen necessary to preserve the transplant, even with caregiver support.” Changed “Review Date” in header to “Date of Last Revision,” and “Date” in the revision log header to “Revision Date.”	11/21	11/21
Annual review completed. References reviewed and updated. Updated description and background with no clinical significance. Updated all contraindications in criteria I.B. “Experimental/investigational” verbiage replaced in criteria IV. statement with descriptive language.	4/22	04/22

References

1. Robertson, RP. Pancreas and islet transplantation in diabetes mellitus. UpToDate. www.uptodate.com. Published August 27, 2020. Accessed January 24, 2022.
2. Leard LE, Holm AM, Valapour M, et al. Consensus document for the selection of lung transplant candidates: An update from the International Society for Heart and Lung Transplantation. *J Heart Lung Transplant*. 2021;40(11):1349-1379. doi:10.1016/j.healun.2021.07.005
3. American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care*. 2014;37 Suppl 1:S81-S90. doi:10.2337/dc14-S081

CLINICAL POLICY **Pancreas Transplantation**

4. Centers for Disease Control and Prevention. National Diabetes Statistics Report, 2020. Atlanta, GA: Centers for Disease Control and Prevention, US Dept of Health and Human Services; 2020.
5. Atkinson MA, Eisenbarth GS, Michels AW. Type 1 diabetes. *Lancet*. 2014;383(9911):69-82. doi:10.1016/S0140-6736(13)60591-7
6. Chiang JL, Kirkman MS, Laffel LM, Peters AL; Type 1 Diabetes Sourcebook Authors. Type 1 diabetes through the life span: a position statement of the American Diabetes Association. *Diabetes Care*. 2014;37(7):2034-2054. doi:10.2337/dc14-1140
7. Mittal S, Johnson P, Friend P. Pancreas transplantation: solid organ and islet. *Cold Spring Harb Perspect Med*. 2014;4(4):a015610. Published 2014 Apr 1. doi:10.1101/cshperspect.a015610
8. Alhamed T, Stratta RJ. Pancreas-kidney transplantation in diabetes mellitus: Patient selection and pretransplant evaluation. UpToDate. www.uptodate.com. Published October 05, 2021. Accessed January 25, 2022.
9. Alhamed T, Stratta RJ. Pancreas-kidney transplantation in diabetes mellitus: Benefits and complication. UpToDate. www.uptodate.com. Published October 05, 2021. Accessed January 24, 2022.
10. Gruessner AC. 2011 update on pancreas transplantation: comprehensive trend analysis of 25,000 cases followed up over the course of twenty-four years at the International Pancreas Transplant Registry (IPTR). *Rev Diabet Stud*. 2011;8(1):6-16. doi:10.1900/RDS.2011.8.6
11. Robertson RP, Davis C, Larsen J, Stratta R, Sutherland DE. Pancreas and islet transplantation for patients with diabetes. *Diabetes Care*. 2000;23(1):112-116. doi:10.2337/diacare.23.1.112
12. Redfield RR, Scalea JR, Odorico JS. Simultaneous pancreas and kidney transplantation: current trends and future directions. *Curr Opin Organ Transplant*. 2015;20(1):94-102. doi:10.1097/MOT.0000000000000146
13. Kesseli SJ, Smith KA, Gardner TB. Total pancreatectomy with islet autologous transplantation: the cure for chronic pancreatitis? *Clin Transl Gastroenterol*. 2015;6(1):e73. Published 2015 Jan 29. doi:10.1038/ctg.2015.2

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. "Health Plan" means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan's affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

CLINICAL POLICY Pancreas Transplantation

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note: For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

Note: For Medicare members, to ensure consistency with the Medicare National Coverage Determinations (NCD) and Local Coverage Determinations (LCD), all applicable NCDs, LCD's and Medicare Coverage Articles should be reviewed prior to applying the criteria set forth in this clinical policy. Refer to the CMS website at <http://www.cms.gov> for additional information.

©2016 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.